

# Effects of Age and Military Service on Strength and Physiological Characteristics of U.S. Army Soldiers

John P. Abt, PhD\*; Katherine Perlsweig, MS†; Takashi Nagai, PhD‡; Timothy C. Sell, PhD‡;  
COL Michael D. Wirt, MC USA‡; Scott M. Lephart, PhD\*

**ABSTRACT** Soldiers must maintain tactical performance capabilities over the course of their career. Loss in physical readiness may be a function of age and the operational demands associated with increasing years of service. The purpose of this study was to assess strength and physiological characteristics in different cohorts of U.S. Army Soldiers based on years of service and age. A total of 253 Soldiers (age:  $28.1 \pm 6.8$  years; height:  $1.76 \pm 0.11$  m; mass:  $84.1 \pm 12.2$  kg) participated. Individual subject cohorts were created based on years of service (1–5 years, 6–10 years, 11–15 years) and age (20–24 years, 25–29 years, 30–34 years, 35–39 years, 40–44 years). Testing included shoulder, knee, ankle, and torso strength, aerobic capacity/lactate threshold, anaerobic power/capacity, and body composition/total mass. Those with 11 to 15 years of service and between ages 30 and 34 had a higher percentage of body fat, and lower aerobic capacity and lactate threshold than younger Soldiers with fewer years of service. Physical training interventions should focus on maintenance of physiological characteristics to offset the loss of readiness at the similar time point of 11 to 15 years of service and 30 to 34 years of age.

## INTRODUCTION

A career in the armed forces is demanding given the high operational tempo and wide spectrum of mission requirements. Service members must maintain a high level of physical readiness to meet the operational demands.<sup>1,2</sup> Although service members are a diverse cohort,<sup>3–5</sup> those with the same military occupational specialty are required to complete similar training and operational tasks despite differences in age and years of military service.<sup>6–8</sup>

Changes in aerobic fitness because of age have been well documented in the civilian population. As age increases, maximum heart rate decreases,<sup>9–12</sup> reducing maximum cardiac output during exercise.<sup>13</sup> The resulting decline in maximal oxygen consumption<sup>9,11,14</sup> is nonlinear,<sup>15,16</sup> with an accelerated deterioration after 45 to 50 years old.<sup>15,17</sup> Power and strength decreases are also nonlinear,<sup>18,19</sup> with power decreasing by 3% per decade from ages 24 to 50 and a 7% decline per decade from ages 50 to 74.<sup>20</sup> Brown, et al<sup>14</sup> measured a 0.048 watt per kilogram decline per year when exercising at maximum aerobic capacity, demonstrating the decline in power produced at maximal effort as age increases.

Ageing results in decreased cardiorespiratory fitness and increased body fat,<sup>16</sup> and these deficits are of particular concern to service members in physically demanding military occupational specialty that are compounded by other

external factors. The weight added by basic protective equipment alone is enough to decrease maximum aerobic capacity by 50% and increase overall caloric expenditure by 20%.<sup>21</sup> Previous research has also found decreased aerobic fitness after deployment<sup>22–24</sup> and decreased anaerobic power after as little as 72 hours of sustained operational training.<sup>25,26</sup> Deployment has been associated with increased body mass, fat mass, and percentage of body fat,<sup>22</sup> with small strength gains<sup>22,24</sup> as Soldiers self-select strength exercises more frequently than aerobic conditioning.<sup>23,24,27</sup> Although these changes because of deployment and training are relatively small, they may be amplified by age-related physiological changes. If there is insufficient dwell time to rest, recover, and retrain between deployments and sustained operational training, these effects may become permanent.<sup>23</sup>

Strength and physiological characteristics such as aerobic and anaerobic capacity are essential to performance of military tasks. The relationship between these characteristics, age, and years of military service are unknown. The purpose of this study was to assess strength and physiological characteristics in different cohorts of U.S. Army Soldiers based on years of service and age. We hypothesized that older Soldiers and those with more years of service would have a poorer physical and physiological profiles than their younger, less experienced counterparts.

## METHODS

### PARTICIPANTS

A total of 253 male Soldiers from the Army 101st Airborne Division (Air Assault) were recruited to participate in this study (age:  $28.1 \pm 6.8$  years; height:  $1.76 \pm 0.11$  m; mass:  $84.1 \pm 12.2$  kg). Human subject's approval was obtained from the respective civilian and military Institutional Review Boards.

\*College of Health Sciences, University of Kentucky, UK Wethington Building, Room 123, 900 South Limestone, Lexington, KY 40536.

†Neuromuscular Research Laboratory, Warrior Human Performance Research Center, University of Pittsburgh, 3830 South Water Street, Pittsburgh, PA 15203.

‡US Army Institute of Surgical Research, 3698 Chambers Pass, Building 3611, JBSA Fort Sam Houston, TX 78234-6315.

doi: 10.7205/MILMED-D-15-00036

## Study Design

This study is a cross-sectional cohort design comparing groups of Soldiers based on years of service (1–5 years, 6–10 years, 11–15 years) and age (20–24 years, 25–29 years, 30–34 years, 35–39 years, 40–44 years). All testing was performed at the University of Pittsburgh Human Performance Research Center at Fort Campbell, Kentucky. Testing was performed on two separate days, with at least 24 hours between test days.

## Instrumentation and Procedures

Body composition (percent body fat: %BF) was measured using a BOD POD Body Composition System (Cosmed, Italy). The BOD POD uses air displacement plethysmography to measure body volume and calculate body density. This has been shown to be a reliable (intraclass correlation coefficient [ICC] = 0.98–0.996) and valid (standard error of measurement [SEM] = 0.47% BF) method of measuring body composition<sup>28</sup> in heterogeneous samples across a variety of populations.<sup>29–31</sup> The system was calibrated using the manufacturer recommended two-point calibration before each test. Subjects were tested according to the manufacturer's protocol and as previously reported.<sup>32</sup>

Bilateral isokinetic strength of the knee flexors/extensors, shoulder internal/external rotators, and torso rotators were measured using the Biodex Multi-Joint System 3 Pro (Biodex Medical Systems, Shirley, New York). Isokinetic dynamometry has been shown to be highly reliable (knee flexion: ICC = 0.93–0.98, knee extension: ICC = 0.96–0.97, shoulder internal and external rotation: ICC = 0.784–0.798, trunk rotation: ICC = 0.89–0.906) and a valid measure of the strength of the primary movers in each of these ranges of motion.<sup>28,33–36</sup> Subjects were positioned and stabilized according to the manufacturer's guidelines, ensuring proper alignment for testing and restricting accessory movements. The subject then performed three practice trials at 50% maximal effort and three warm-up trials at maximal effort, followed by 1 minute of rest. Peak isokinetic torque was then recorded across five maximal effort repetitions (concentric/concentric at 60° per second) and reported normalized to percent body weight. Strength ratios were calculated to compare agonist–antagonist strength of knee flexion to extension, shoulder external to internal rotation, and left to right torso rotation.

Anaerobic power and capacity were measured using an electronically braked Velotron cycling ergometer (Racermate, Seattle, Washington) during a Wingate protocol. This has been shown to be a valid.<sup>37,38</sup> and reliable (ICC = 0.91) measure of an individual's ability to perform short-term, high intensity exercise.<sup>28</sup> The Velotron was calibrated according to factory recommendations, and the seat and handle bars were adjusted to fit each subject. After warming up at a self-selected pace, subjects pedaled at 125 W for 20 seconds, and then performed a maximal effort sprint for 30 seconds against a braking torque of 9% body weight. This protocol has been

previously reported.<sup>28</sup> Anaerobic power was reported as the peak watts normalized to body weight during the first 5 seconds of the test, and anaerobic capacity was reported as the average watts normalized to body weight produced during the entire 30 seconds.

Aerobic capacity ( $VO_{2max}$ ) and lactate threshold were measured during an incremental treadmill protocol using a portable metabolic system (Oxycon Mobile; Viasys, San Francisco, California), heart rate monitor (Polar USA, Lake Success, New York), and lactate analyzer (Arkay, Inc., Kyoto, Japan). These systems are valid measurement tools,<sup>28,39</sup> and were calibrated according to factory specifications before each test. Subjects performed a warm-up at a self-selected pace on the treadmill for 5 minutes before testing. The modified incremental protocol used a constant speed and a 2.5% increase in grade at the end of each 3 minute stage until volitional fatigue.<sup>28,40</sup> Treadmill speed was set at 70% of the mile pace from the subject's most recent Army Physical Fitness Test 2-mile run time.<sup>28</sup> Aerobic capacity was normalized to body weight (mL/kg/min) to evaluate differences in aerobic fitness between subjects. Blood lactate levels were collected via finger stick at rest, in the last minute of the warm-up and each 3 minute test stage, immediately upon test termination, and 3 minutes after test completion. Lactate threshold (the inflection point where blood lactate increased nonlinearly) was reported in relation to absolute value of oxygen consumption ( $VO_2$ ), and the relative values of percentage of  $VO_{2max}$  and heart rate at lactate threshold.

## Statistical Analysis

Shapiro-Wilk and Levene's tests were used to determine normality of sample distribution and homogeneity of variance respectively. For normally distributed variables, one-way analysis of variance (ANOVA) and Tukey's post-hoc procedures were used to determine differences between age and years of service groups. For variables with significant Shapiro-Wilk tests, a Kruskal-Wallis ANOVA was used to determine differences between groups. For those with significant Levene's tests, an ANOVA with planned comparisons and adjusted *p* values were used. All statistical measures were obtained using IBM SPSS Statistics for Windows (Version 21, IBM Corp., Armonk, NY). Statistical significance was set at *p* < 0.05.

## RESULTS

Descriptive data for each age cohort are presented in Table I. There were no significant differences for height or mass between age cohorts (*p* > 0.05). Physiological data for the age cohort are presented in Table II. Soldiers aged 30 to 34 had more body fat than Soldiers aged 20 to 24 (*p* = 0.005) and 25 to 29 (*p* = 0.012). Aerobic capacity ( $VO_{2max}$ ) was higher in 20 to 24 year olds than 30 to 34 year olds (*p* = 0.041), 35 to 39 year olds (*p* = 0.047) and 40 to 44 year olds (*p* = 0.041). Although there was no change in the percentage

TABLE I. Descriptive Statistics by Age

Age	n	Age (Years)		Experience (Years)		Height (cm)*		Weight (kg) No.	
		Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
20–24	77	21.13	1.35	3.83	1.36	176.98	7.55	80.82	12.10
25–29	79	26.61	1.29	5.96	2.54	178.12	7.13	86.11	11.43
30–34	45	31.53	1.50	9.69	2.63	177.4	6.31	85.43	13.70
35–39	19	36.79	1.62	10.58	3.34	176.2	7.21	86.33	8.22
40–44	21	42.1	1.76	12.81	1.57	175.15	9.33	83.34	12.54

\*No significant differences between age cohorts ( $p > 0.05$ ).

of  $VO_{2max}$  at which lactate threshold occurred, the absolute value of  $VO_2$  at lactate threshold decreased with age:  $VO_2$  at lactate threshold was higher in 25 to 29 year olds than 30 to 35 year olds ( $p = 0.0207$ ). Heart rate at lactate threshold was significantly greater in those age 20 to 24 than those age 40 to 44 ( $p = 0.035$ ). The overall analysis for anaerobic capacity was significant ( $p = 0.01$ ), but there were no significant post-hoc comparisons. No significant strength differences existed for knee flexion, knee extension, shoulder internal rotation, shoulder external rotation, torso rotation, right knee flexion/

extension ratio, left and right shoulder external rotation/internal rotation ratio, or torso rotation ratio ( $p > 0.05$ ). The overall analysis for left knee flexion/extension ratio was significant ( $p = 0.035$ ), but there were no significant post-hoc comparisons.

Descriptive data for each years of service cohort are presented in Table III. There were no significant differences for height or mass between the years of service cohorts ( $p > 0.05$ ). For physiological data (Table IV), Soldiers with 1 to 5 years of experience had lower body mass index (BMI) than those with 11 to 15 years of service ( $p = 0.023$ ).

TABLE II. Results Summary by Age

	20–24		25–29		30–34		35–39		40–44	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
<b>Physiological Data</b>										
Body Fat (%) <sup>a</sup>	18.79	7.26	19.26	7.55	23.32	7.67	24.52	5.31	21.61	5.63
BMI	25.82	3.48	27.38	3.62	27.42	3.93	27.82	2.02	27.82	2.51
HR <sub>max</sub> (bpm)	191.99	9.05	187.35	9.18	188.43	7.03	184.41	8.64	179.42	8
VO <sub>2max</sub> (mL/kg/min) <sup>b</sup>	48.73	6.33	48.07	7.22	45.11	6.41	43.65	4.76	43.07	6.88
VO <sub>2</sub> at LT (mL/kg/min) <sup>c</sup>	39.06	6.28	40.07	6.61	36.35	5.56	36.23	4.38	37.09	6.23
LT (%VO <sub>2max</sub> )	80.33	9.04	83.37	6.53	80.82	7.11	83.17	7.73	86.25	6.56
HR at LT (bpm) <sup>d</sup>	170.86	12.82	170.04	11.55	166.78	9.34	165.76	9.6	164.63	8.31
%HR <sub>max</sub> at LT	89.01	5.39	90.76	4.38	88.55	4.39	89.94	4.38	91.81	3.74
Anaerobic Power (w/kg)	13.29	1.92	14.07	2.13	13.43	1.89	13.05	2.19	12.99	1.51
Anaerobic Capacity (w/kg)	7.84	0.93	7.97	1.17	7.57	0.95	7.43	0.82	7.46	0.92
<b>Strength Data*</b>										
Left Knee Flexion (%BW)	113.06	27.75	114.52	26.11	115.84	22.75	109.79	21.94	109.49	27.53
Right KF (%BW)	115.49	26.66	120.52	26.36	112.87	26.62	116.15	20.92	115.09	22.57
Left KE (%BW)	229.81	43.67	228.74	45.36	231.62	43.8	208.75	34.28	204.04	46.78
Right KE (%BW)	241.67	48.92	238.68	49.37	243.41	45.92	223.93	40.78	224.58	44.19
Left KF:KE Ratio	0.49	0.08	0.5	0.09	0.51	0.08	0.53	0.09	0.55	0.12
Right KF:KE Ratio	0.48	0.08	0.51	0.08	0.47	0.09	0.53	0.08	0.52	0.07
Left Shoulder IR (%BW)	54.61	16.02	55.63	15.22	56.64	15.14	53.14	13.36	57.82	15.84
Right Shoulder IR (%BW)	60.82	14.95	61.35	14.4	62.79	14.8	57	12.93	59.18	17.1
Left Shoulder ER (%BW)	38.61	7.21	37.54	8.07	37.35	8.29	35.86	6.31	35.99	7.56
Right Shoulder ER (%BW)	42.99	8.61	41.69	8.69	41.89	8.62	39.35	7.26	38.49	7.7
Left Shoulder ER:IR Ratio	0.74	0.16	0.7	0.15	0.68	0.11	0.69	0.12	0.64	0.12
Right Shoulder ER:IR Ratio	0.73	0.16	0.69	0.12	0.68	0.1	0.7	0.1	0.67	0.12
Left TR (%BW)	150.18	37.47	157.75	29.56	154.15	31.99	144.52	25.79	153.65	38.35
Right TR (%BW)	149.99	34.43	156.21	30.26	155.57	33.15	144.87	22.13	153.69	36.04
Left:Right TR Ratio	1.01	0.12	1.02	0.12	1	0.11	1	0.11	1	0.1

BMI, body mass index;  $VO_2$ , aerobic capacity;  $VO_{2max}$ , maximum aerobic capacity; LT, lactate threshold; HR, heart rate in beats per minute; HR<sub>max</sub>, maximum heart rate; KE, knee extension; KF, knee flexion; IR, internal rotation; ER, external rotation; TR, trunk rotation. Physiology data. <sup>a</sup>Body fat higher in Soldiers aged 30 to 34 compared to 20 to 24 ( $p = 0.005$ ) and 25 to 29 ( $p = 0.012$ ). <sup>b</sup>Aerobic capacity ( $VO_{2max}$ ) was higher in 20 to 24 year olds than 30 to 34 year olds ( $p = 0.041$ ), 35 to 39 year olds ( $p = 0.047$ ) and 40 to 44 year olds ( $p = 0.041$ ). <sup>c</sup> $VO_2$  at LT was higher in 25 to 29 year olds than 30 to 35 year olds ( $p = 0.0207$ ). <sup>d</sup>Heart rate at LT was significantly greater in those age 20 to 24 than those age 40 to 44 ( $p = 0.035$ ) Strength data. \*No significant differences between age cohorts for knee, shoulder, or torso strength ( $p > 0.05$ ).

**TABLE III.** Descriptive Statistics by Years of Experience

Years	n	Experience (years)		Age (years)		Height (cm)*		Weight (kg)*	
		Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
1–5	123	3.53	1.08	23.68	3.42	177.36	7.60	82.95	12.88
6–10	67	7.63	1.23	28.46	3.89	177.02	7.62	84.91	10.88
11–15	63	12.49	0.95	36.51	5.77	176.49	7.79	85.49	11.99

\*No significant differences between age cohorts ( $p > 0.05$ ).

Soldiers with 1 to 5 ( $p < 0.001$ ) and 6 to 10 years ( $p = 0.016$ ) of service both had significantly less body fat than those with 11 to 15 years of service. Soldiers with 1 to 5 ( $p < 0.001$ ) and 6 to 10 ( $p < 0.001$ ) years of service had higher  $VO_{2max}$  than those with 11 to 15 years of service. Lactate threshold was also higher in those with 1 to 5 years of service compared to the 11 to 15 years of service group ( $p = 0.017$ ). Heart rate at lactate threshold was significantly

greater in those with 1 to 5 years of service than those with 11 to 15 years of service ( $p = 0.012$ ). Left knee flexion/extension ratio was significantly lower in the 1 to 5 years of service compared to the 6 to 10 years of service group, and both the 1 to 5 and 6 to 10 years of experience groups had lower ratios than the 11 to 15 years of service group ( $p < 0.001$  for all three comparisons). Left and right shoulder internal/external rotation strength ratios were significantly

**TABLE IV.** Results Summary by Years of Experience

	1–5		6–10		11–15	
	Mean	±SD	Mean	±SD	Mean	±SD
<b>Physiological Data</b>						
Body Fat (%) <sup>a</sup>	19.11	7.43	20.48	7.72	23.25	5.73
BMI <sup>b</sup>	26.48	3.82	27.08	3.50	27.86	2.83
HR <sub>max</sub> (bpm)	190.54	9.16	186.77	8.33	185.35	9.89
$VO_{2max}$ (mL/kg/min) <sup>c</sup>	48.37	6.94	47.31	6.85	44.04	5.69
$VO_2$ at LT <sup>d</sup>	39.45	6.53	38.76	6.30	36.79	5.22
% $VO_{2max}$ at LT	81.71	8.10	82.01	7.24	83.73	7.56
HR at LT <sup>e</sup>	171.15	11.35	167.32	11.85	166.78	9.61
%HR <sub>max</sub> at LT	89.85	4.73	89.58	4.82	90.08	4.59
Anaerobic Power (w/kg)	13.42	1.80	13.69	2.12	13.49	2.14
Anaerobic Capacity (w/kg)	7.84	0.92	7.79	1.15	7.62	1.03
<b>Strength Data</b>						
Left KF (%BW)	112.92	27.03	113.31	24.18	113.57	24.50
Right KF (%BW)	116.54	27.44	114.56	23.27	117.06	24.55
Left KE (%BW)	230.17	45.92	224.59	39.41	215.97	45.07
Right KE (%BW)	244.44	52.04	230.78	39.60	233.53	43.07
Left KF:KE Ratio*	0.49	0.08	0.51	0.08	0.53	0.10
Right KF:KE Ratio	0.48	0.08	0.50	0.08	0.51	0.08
Left Shoulder IR (%BW)	54.60	16.07	58.01	14.01	54.39	13.95
Right Shoulder IR (%BW)	60.18	15.32	63.83	13.99	58.98	13.33
Left Shoulder ER (%BW)	38.30	7.86	37.29	7.70	36.09	6.61
Right Shoulder ER (%BW)	42.43	9.12	41.62	7.85	39.85	7.59
Left Shoulder ER:IR Ratio#	0.73	0.15	0.66	0.14	0.68	0.01
Right Shoulder ER:IR Ratio#	0.73	0.14	0.66	0.11	0.69	0.11
Left TR (%BW)	151.32	33.57	157.97	35.35	150.86	31.19
Right TR (%BW)	152.20	31.48	154.14	35.94	152.02	29.75
Left:Right TR Ratio	1.00	0.11	1.04	0.13	1.00	0.11

BMI, body mass index;  $VO_2$ , aerobic capacity;  $VO_{2max}$ , maximum aerobic capacity; LT, lactate threshold; HR, heart rate in beats per minute; HR<sub>max</sub>, maximum heart rate; KE, knee extension; KF, knee flexion; IR, internal rotation; ER, external rotation; TR, trunk rotation. Physiology Data. <sup>a</sup>Soldiers with 1 to 5 ( $p < 0.001$ ) and 6 to 10 years ( $p = 0.016$ ) of service both had significantly less body fat than those with 11 to 15 years of service. <sup>b</sup>Soldiers with 1 to 5 years of experience had lower BMI than those with 11 to 15 years of service ( $p = 0.023$ ). <sup>c</sup>Soldiers with 1 to 5 ( $p < 0.001$ ) and 6 to 10 ( $p < 0.001$ ) years of service had higher  $VO_{2max}$  than those with 11 to 15 years of service. <sup>d</sup>LT was higher in those with 1 to 5 years of service compared to the 11 to 15 years of service group ( $p = 0.017$ ). <sup>e</sup>Heart rate at LT was significantly greater in those with 1 to 5 years of service than those with 11 to 15 years of service ( $p = 0.012$ ) Strength Data. \*Left knee flexion/extension ratio was significantly lower in the 1 to 5 years of service compared to the 6 to 10 years of service group, and both the 1 to 5 and 6 to 10 years of experience groups had lower ratios than the 11 to 15 years of service group ( $p < 0.001$  for all three comparisons) #Left and right shoulder internal/external rotation strength ratios were significantly greater in those with 1 to 5 years of experience compared to those with 6 to 10 years of experience (Left:  $p = 0.001$ , Right:  $p = 0.007$ ).

greater in those with 1 to 5 years of experience compared to those with 6 to 10 years of experience (left:  $p = 0.001$ , right:  $p = 0.007$ ). There were no significant strength differences for knee flexion, knee extension, shoulder internal rotation, shoulder external rotation, torso rotation, or torso rotation ratio ( $p > 0.05$ ).

## DISCUSSION

Older Soldiers and those with more years of service demonstrated poorer physiological characteristics, with large declines occurring at the similar time period of age 30 to 34 and 11 to 15 years of service. Most Soldiers enlist in the Army between ages 19 and 21,<sup>5</sup> meaning that a 30-year-old Soldier likely has 11 years of service. Those with 11 to 15 years of service and between ages 30 and 34 had a higher percentage of body fat, and lower maximum heart rate, aerobic capacity, and lactate threshold than younger soldiers with fewer years of service. Strength ratios at the knee and shoulder were significantly different across the experience groups, indicating that strength ratios may be more sensitive to change than discrete strength measures.

Although the Department of Defense's goal is to allow a service member twice the amount of recovery time as deployment time, the ratios of deployment to dwell time during Operation Iraqi Freedom and Operation Enduring Freedom were closer to 1:1.<sup>41</sup> This lack of recovery and retraining time may explain why older soldiers and those with more years of service had a larger decrement in performance variables than would be expected in a civilian or younger population. Changes in aerobic capacity in civilian athletes have been attributed largely to declines in  $HR_{max}$ ,<sup>42</sup> which decreases at approximately 0.7 to 1 beat/min/year.<sup>10,12,43</sup> This steady decline is similar to that observed between age and experience groups in the present study. Astrand's classic study of maximal oxygen uptake and age reported a 7% decline in  $HR_{max}$  over the 20-year observation period<sup>9</sup>; nearly identical to the 6.7% decrease in  $HR_{max}$  between soldiers age 20 to 24 and those age 40 to 44.

Aerobic capacity decreases in a nonlinear pattern among civilian athletes, with large decreases in  $VO_{2max}$  measured after age 45 to 50,<sup>15,17,20,44</sup> unlike in this sample, where a large decrease occurred at age 30 to 34. These results are similar to those of Giovannetti, et al<sup>45</sup>, who observed no significant change in estimated  $VO_{2max}$  until age 30 to 39 in members of the U.S. Air Force; with a large drop in aerobic capacity after age 30 and a steady decline continuing through age 50 and over. This decline in  $VO_{2max}$  may be due in part to increased body fat in older and more experienced Soldiers, as previous research has shown that men who are able to maintain lean body mass are more likely to see smaller declines in aerobic capacity<sup>42</sup> and cardiorespiratory fitness.<sup>15</sup> The absolute declines in  $HR_{max}$  and  $VO_{2max}$  observed here resulted in lower heart rate and  $VO_2$  at lactate threshold: with no significant difference in percentage of maximum heart rate or percentage of  $VO_{2max}$  at lactate threshold, lower maximal

values for these variables mean that the absolute values at lactate threshold must be lower as well.

Despite similarities in age-related changes in  $HR_{max}$  and body composition between civilian athletes and Soldiers, the combined effects of operational stress and deployment likely result in greater decrements in aerobic capacity in this cohort. Soldiers with fitness deficiencies must exert a larger percentage of their maximum effort when performing work related tasks, accelerating fatigue and potentially increasing risk for injury.<sup>28,46,47</sup> Studies of injury rates in Soldiers identify decreased aerobic capacity,<sup>3,7,24,45-49</sup> higher BMI<sup>4,48,50</sup> and increased percentage of body fat<sup>28,49</sup> as risk factors for injury development. Therefore, the decrease in aerobic capacity and increase in body fat observed in Soldiers of 30 to 34 years of age and with 11 to 15 years of service in this study are especially concerning for risk of injury.

Although it was statistically significant, left knee flexion to extension ratio only increased by 0.04 between the 1 to 5 and 11 to 15 years of service groups, and this measure was approximately 0.50 for all age and experience groups on the left and right knees. This is well below the 0.60 hamstring to quadriceps strength ratio recommended for ligamentous and muscular injury prevention.<sup>51</sup> Soldiers from all age and experience groups could benefit from a hamstring strengthening program. Bilateral shoulder internal to external rotation strength ratios were statistically higher in soldiers with 1 to 5 years of service compared to those with 6 to 10 years of service; however, the means for both of these groups (left:  $0.73 \pm 0.14$  vs.  $0.66 \pm 0.14$ ; right:  $0.72 \pm 0.14$  vs.  $0.66 \pm 0.11$ ) are within the normative range identified by Ellenbecker et al<sup>52</sup> of 0.66 to 0.75.

## CONCLUSION

Changes in strength and physiological characteristics occurred at younger ages in this cohort of Army personnel than has been observed in the civilian population. Although the exact reasons for the early decline in aerobic capacity and increased body fat are not completely clear, they may be because of the cumulative effects of operational training and deployment. Strength findings in this study suggest that muscle balance is more sensitive to small changes than individual muscle measures; therefore, human performance staff should use strength ratios to monitor possible strength decrements across a Soldier's tactical lifecycle. Physical training interventions for force protection have been effective,<sup>53-56</sup> and the results of this study indicate that these efforts should focus on Soldiers age 30 to 34, particularly for those with more years of service.

## ACKNOWLEDGMENT

This work was supported by the U.S. Army Medical Research and Materiel Command No. W81XWH-11-2-0097. Opinions, interpretations, conclusions, and recommendations are those of the authors and not necessarily endorsed by the Department of Defense or U.S. Army.

## REFERENCES

- Skeehan CD, Tribble DR, Sanders JW, Putnam SD, Armstrong AW, Riddle MS: Nonbattle injury among deployed troops: an epidemiologic study. *Mil Med* 2009; 174(12): 1256–62.
- U.S. Department of the Army: Army Physical Readiness Training. Field Manual 7–22. Washington, DC: Headquarters, Department of the Army, May 13, 2013. Available at [http://armypubs.army.mil/doctrine/DR\\_pubs/dr\\_a/pdf/fm7\\_22.pdf](http://armypubs.army.mil/doctrine/DR_pubs/dr_a/pdf/fm7_22.pdf); accessed October 14, 2014.
- Knapik J, Ang P, Reynolds K, Jones B: Physical fitness, age, and injury incidence in infantry soldiers. *J Occup Med* 1993; 35(6): 598–603.
- Packnett ER, Niebuhr DW, Bedno SA, Cowan DN: Body mass index, medical qualification status, and discharge during the first year of US Army service. *Am J Clin Nutr* 2011; 93(3): 608–14.
- Sharp MA, Patton JF, Knapik JJ, et al: Comparison of the physical fitness of men and women entering the U.S. Army: 1978–1998. *Med Sci Sports Exerc* 2002; 34(2): 356–63.
- Henning PC, Park BS, Kim JS: Physiological decrements during sustained military operational stress. *Mil Med* 2011; 176(9): 991–7.
- Jones BH, Cowan DN, Tomlinson JP, Robinson JR, Polly DW, Frykman PN: Epidemiology of injuries associated with physical training among young men in the army. *Med Sci Sports Exerc* 1993; 25(2): 197–203.
- Friedl KE: Body composition and military performance—many things to many people. *J Strength Cond Res* 2012; 26(Suppl 2): S87–100.
- Astrand I, Astrand PO, Hallbäck I, Kilbom A: Reduction in maximal oxygen uptake with age. *J Appl Physiol* 1973; 35(5): 649–54.
- Katzel LI, Sorkin JD, Fleg JL: A comparison of longitudinal changes in aerobic fitness in older endurance athletes and sedentary men. *J Am Geriatr Soc* 2001; 49(12): 1657–64.
- Kusy K, Zielinski J: Aerobic capacity in speed–power athletes aged 20–90 years vs endurance runners and untrained participants. *Scand J Med Sci Sports* 2014; 24(1): 68–79.
- Tanaka H, Monahan KD, Seals DR: Age-predicted maximal heart rate revisited. *J Am Coll Cardiol* 2001; 37(1): 153–6.
- Tanaka H, Seals DR: Endurance exercise performance in Masters athletes: age-associated changes and underlying physiological mechanisms. *J Physiol* 2008; 586(1): 55–63.
- Brown SJ, Ryan HJ, Brown JA: Age-associated changes in  $\text{VO}_2$  and power output—A cross-sectional study of endurance trained New Zealand Cyclists. *J Sports Sci Med* 2007; 6(4): 477–83.
- Jackson AS, Sui X, Hébert JR, Church TS, Blair SN: Role of lifestyle and aging on the longitudinal change in cardiorespiratory fitness. *Arch Intern Med* 2009; 169(19): 1781–7.
- Wiswell RA, Hawkins SA, Jaque SV, et al: Relationship between physiological loss, performance decrement, and age in master athletes. *J Gerontol A Biol Sci Med Sci* 2001; 56(10): M618–26.
- Pimentel AE, Gentile CL, Tanaka H, Seals DR, Gates PE: Greater rate of decline in maximal aerobic capacity with age in endurance-trained than in sedentary men. *J Appl Physiol* (1985); 2003; 94(6): 2406–13.
- Kasch FW, Boyer JL, Schmidt PK, et al: Ageing of the cardiovascular system during 33 years of aerobic exercise. *Age Ageing* 1999; 28(6): 531–6.
- Metter EJ, Lynch N, Conwit R, Lindle R, Tobin J, Hurley B: Muscle quality and age: cross-sectional and longitudinal comparisons. *J Gerontol A Biol Sci Med Sci* 1999; 54(5): B207–18.
- Seiler KS, Spirduso WW, Martin JC: Gender differences in rowing performance and power with aging. *Med Sci Sports Exerc* 1998; 30(1): 121–7.
- Sell TC, Chu Y, Abt JP: Minimal additional weight of combat equipment alters air assault soldiers' landing biomechanics. *Mil Med* 2010; 175(1): 41–7.
- Lester ME, Knapik JJ, Catrambone D, et al: Effect of a 13-month deployment to Iraq on physical fitness and body composition. *Mil Med* 2010; 175(6): 417–23.
- Sharp MA, Knapik JJ, Walker LA, et al: Physical fitness and body composition after a 9-month deployment to Afghanistan. *Med Sci Sports Exerc* 2008; 40(9): 1687–92.
- Warr BJ, Heumann KJ, Dodd DJ, Swan PD, Alvar BA: Injuries, changes in fitness, and medical demands in deployed National Guard soldiers. *Mil Med* 2012; 177(10): 1136–42.
- Hackney AC, Shaw JM, Hodgdon JA, Coyne JT, Kelleher DL: Cold exposure during military operations: effects on anaerobic performance. *J Appl Physiol* (1985) 1991; 71(1): 125–30.
- Nindl BC, Leone CD, Tharion WJ, et al: Physical performance responses during 72 h of military operational stress. *Med Sci Sports Exerc* 2002; 34(11): 1814–22.
- Nindl BC, Castellani JW, Warr BJ, et al: Physiological Employment Standards III: physiological challenges and consequences encountered during international military deployments. *Eur J Appl Physiol* 2013; 113(11): 2655–72.
- Crawford K, Fleishman K, Abt JP, Sell TC, et al: Less body fat improves physical and physiological performance in army soldiers. *Mil Med* 2011; 176(1): 35–43.
- Ballard TP, Fafara L, Vukovich MD: Comparison of Bod Pod and DXA in female collegiate athletes. *Med Sci Sports Exerc* 2004; 36(4): 731–5.
- Noreen EE, Lemon PW: Reliability of air displacement plethysmography in a large, heterogeneous sample. *Med Sci Sports Exerc* 2006; 38(8): 1505–9.
- Vescovi JD, Zimmerman SL, Miller WC, Hildebrandt L, Hammer RL, Fernhall B: Evaluation of the BOD POD for estimating percentage body fat in a heterogeneous group of adult humans. *Eur J Appl Physiol* 2001; 85(3–4): 326–32.
- Siri WE: Body composition from fluid spaces and density: analysis of methods. In: *Techniques for measuring body composition*, pp 223–4. Edited by Brozek J, Henchel A. Washington, DC, National Academic Sciences/National Research Council, 1961. Available at [http://books.google.com/books/about/Techniques\\_for\\_Measuring\\_Body\\_Compositio.html?id=ATErAAAAAYAAJ](http://books.google.com/books/about/Techniques_for_Measuring_Body_Compositio.html?id=ATErAAAAAYAAJ); accessed October 14, 2014.
- Keskula DR, Dowling JS, Davis VL, Finley PW, Dell'omo DL: Inter-rater reliability of isokinetic measures of knee flexion and extension. *J Athl Train* 1995; 30(2): 167–70.
- Sell TC, Tsai YS, Smoliga JM, Myers JB, Lephart SM: Strength, flexibility, and balance characteristics of highly proficient golfers. *J Strength Cond Res* 2007; 21(4): 1166–71.
- Drouin JM, Valovich-mcLeod TC, Shultz SJ, Gansneder BM, Perrin DH: Reliability and validity of the Biodex system 3 pro isokinetic dynamometer velocity, torque and position measurements. *Eur J Appl Physiol* 2004; 91(1): 22–9.
- Sole G, Hamrén J, Milosavljevic S, Nicholson H, Sullivan SJ: Test-retest reliability of isokinetic knee extension and flexion. *Arch Phys Med Rehabil* 2007; 88(5): 626–31.
- Bar-Or O: The Wingate anaerobic test. An update on methodology, reliability and validity. *Sports Med* 1987; 4(6): 381–94.
- Patton JF, Duggan A: An evaluation of tests of anaerobic power. *Aviat Space Environ Med* 1987; 58(3): 237–42.
- Attinger A, Tüller C, Souren T, Tamm M, Schindler C, Brutsche MH: Feasibility of mobile cardiopulmonary exercise testing. *Swiss Med Wkly* 2006; 136(1–2): 13–8.
- Kang J, Chaloupka EC, Mastrangelo MA, Biren GB, Robertson RJ: Physiological comparisons among three maximal treadmill exercise protocols in trained and untrained individuals. *Eur J Appl Physiol* 2001; 84(4): 291–5.
- Bonds TM, Baiocchi D, McDonald LL: Army deployments of OIF and OEF. RAND Corporation, 2010. Available at [http://www.rand.org/pubs/documented\\_briefings/DB587.html](http://www.rand.org/pubs/documented_briefings/DB587.html); accessed October 14, 2014.
- Hawkins SA, Marcell TJ, Victoria Jaque S, Wiswell RA: A longitudinal assessment of change in  $\text{VO}_{2\text{max}}$  and maximal heart rate in master athletes. *Med Sci Sports Exerc* 2001; 33(10): 1744–50.

43. Nybo L, Schmidt JF, Fritzdorf S, Nordsborg NB: Physiological characteristics of an aging Olympic athlete. *Med Sci Sports Exerc* 2014; 46(11): 2132–8.
  44. Martin JC, Farrar RP, Wagner BM, Spirduso WW: Maximal power across the lifespan. *J Gerontol A Biol Sci Med Sci* 2000; 55(6): M311–6.
  45. Giovannetti JM, Bembem M, Bembem D, Cramer J: Relationship between estimated aerobic fitness and injury rates among active duty at an Air Force base based upon two separate measures of estimated cardiovascular fitness. *Mil Med* 2012; 177(1): 36–40.
  46. Grier TL, Morrison S, Knapik JJ, Canham-Chervak M, Jones BH: Risk factors for injuries in the U.S. Army Ordnance School. *Mil Med* 2011; 176(11): 1292–9.
  47. Knapik JJ, Sharp MA, Canham-Chervak M, Hauret K, Patton JF, Jones BH: Risk factors for training-related injuries among men and women in basic combat training. *Med Sci Sports Exerc* 2001; 33(6): 946–54.
  48. Gardner JW, Kark JA, Karnei K, et al: Risk factors predicting exertional heat illness in male Marine Corps recruits. *Med Sci Sports Exerc* 1996; 28(8): 939–44.
  49. Cowan DN, Bedno SA, Urban N, Yi B, Niebuhr DW: Musculoskeletal injuries among overweight army trainees: incidence and health care utilization. *Occup Med (Lond)* 2011; 61(4): 247–52.
  50. Knapik JJ, Graham B, Cobbs J, Thompson D, Steelman R, Jones BH: A prospective investigation of injury incidence and risk factors among army recruits in combat engineer training. *J Occup Med Toxicol* 2013; 8(1): 5.
  51. Holcomb WR, Rubley MD, Lee HJ, Guadagnoli MA: Effect of hamstring-emphasized resistance training on hamstring:quadriceps strength ratios. *J Strength Cond Res* 2007; 21(1): 41–7.
  52. Ellenbecker T, Davies GJ: The application of isokinetics in testing and rehabilitation of the shoulder complex. *J Athl Train* 2000; 35(3): 338–50.
  53. Bullock SH, Jones BH, Gilchrist J, Marshall SW: Prevention of physical training-related injuries recommendations for the military and other active populations based on expedited systematic reviews. *Am J Prev Med* 2010; 38(1 Suppl): S156–81.
  54. Jones BH, Hansen BC: An armed forces epidemiological board evaluation of injuries in the military. *Am J Prev Med* 2000; 18(3 Suppl): 14–25.
  55. Sell TC, Abt JP, Crawford K, et al: Warrior Model for Human Performance and Injury Prevention: Eagle Tactical Athlete Program (ETAP) Part II. *J Spec Oper Med* 2010; 10(4): 22–33.
  56. Zambraski EJ, Yancosek KE: Prevention and rehabilitation of musculoskeletal injuries during military operations and training. *J Strength Cond Res* 2012; 26(Suppl 2): S101–6.
-